

Original citation:

Elliott, Charles M., Stinner, Björn and Venkataraman, Chandrasekhar. (2012) Modelling cell motility and chemotaxis with evolving surface finite elements. *Journal of The Royal Society Interface*, Vol.9 (No.76). pp. 3027-3044. ISSN 1742-5689

Permanent WRAP url:

<http://wrap.warwick.ac.uk/52273>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes the work of researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

<http://dx.doi.org/10.1098/rsif.2012.0276>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

warwick**publications**wrap
highlight your research

<http://go.warwick.ac.uk/lib-publications>

MODELLING CELL MOTILITY AND CHEMOTAXIS WITH EVOLVING SURFACE FINITE ELEMENTS

CHARLES M. ELLIOTT, BJÖRN STINNER, AND CHANDRASEKHAR VENKATARAMAN

ABSTRACT. We present a mathematical and computational framework for the modelling of cell motility. The cell membrane is represented by an evolving surface, with the movement of the cell determined by the interaction of various forces that act normal to the surface. We consider external forces such as those that may arise due to inhomogeneities in the medium and a pressure that constrains the enclosed volume, as well as internal forces that arise from the reaction of the cells surface to stretching and bending. We also consider a protrusive force associated with a reaction-diffusion system posed on the cell membrane, with cell polarisation modelled by this surface reaction-diffusion system. The computational method is based on an evolving surface finite element method. The general method can account for the large deformations that arise in cell motility and allows the simulation of cell migration in three dimensions. We illustrate applications of the proposed modelling framework and numerical method by reporting on numerical simulations of a model for eukaryotic chemotaxis and a model for the persistent movement of keratocytes in two and three space dimensions. Movies of the simulated cells can be obtained from http://homepages.warwick.ac.uk/~maskae/CV_Warwick/Chemotaxis.html

Keywords: cell motility, chemotaxis, surface finite elements, reaction diffusion systems

1. INTRODUCTION

Modelling the directional motility of cells is of much importance especially due to the central role directed cell migration plays in several biological phenomena such as embryonic development, cancer, tissue development and immune responses [1]. Broadly speaking the motile cycle of a cell consists of the following processes: *polarisation* where the cell develops a front and a back through the redistribution of proteins and lipids within the cell, *protrusion* at the leading edge of the cell pushing the front of the cell outwards and *retraction* of the rear of the cell towards the leading edge [2]. Although the main aspects of the motile cycle appear deceptively simple, as further details are added to the modelling various complexities arise. For example in the case of chemotactic eukaryotic cells, the molecular mechanisms that govern gradient sensing and cell polarisation are still not fully understood [3]. Furthermore, it is difficult to quantify the forces associated with motility and only recently has experimental progress been made in this direction [4, 5]. Direct numerical simulation of cell motility necessitates the consideration of deformable surfaces [6, 7] or multi-phase flow models [8, 9] both of which are computationally challenging. Finally, the deformation of the cell surface is linked to the dynamics of actin and other cell resident proteins and these dynamics must be coupled in a consistent way with the evolution of the cell surface.

In this work we present a mathematical framework for the modelling of cell motility and a numerical method for the simulation of such models. The approach we propose uses ideas of existing $2d$ models but generalises these and extends the modelling to the three-dimensional setting. It consists of partial differential equations, specifically those of reaction-diffusion type, posed on the cell boundary coupled to an evolution law for the cell membrane. Further we discuss the inclusion of external forces and illustrate this with a phenomenological model for the interaction between a cell and obstacles. We present a numerical method, based on evolving triangulated surfaces, that consists of an evolving surface finite element

Date: 18th February 2013.

Address: Mathematics Institute, Zeeman Building, University of Warwick, CV4 7AL. Email: c.venkataraman@warwick.ac.uk.

method [10] for the approximation of the surface partial differential equation and a parameterised finite element method [11] for the approximation of the surface evolution law.

It is our hope that the parametric approach we employ will be more efficient than other standard approaches such as phase field [6] or level set methods [12]. The reasoning behind this statement is that our methodology based on triangulated surfaces formulates the problem in one dimension less than other approaches, in which the equations are discretised in the embedding space. See, for example, [13, 14, 15, 16] and references therein for further discussion.

We consider two specific models for cell motility. Firstly a model for eukaryotic chemotaxis. Aspects of chemotaxis, such as changes in direction due to splitting and biased generation of pseudopods as well as response to a changing chemotactic signal [17], are present in this surface based model. We also present simulations of pseudopod-driven migration of a cell in three-dimensions. We next consider a surface based model for the persistent migration of fish keratocytes, presenting numerical simulations in two- and three-dimensions. The surface based model is able to capture the different shapes that characterise migrating keratocytes and the correlation between aspect ratio and cell speed [18].

A summary of the contributions of our study is as follows. We derive a rigorous mathematical framework for the modelling of cell motility and chemotaxis in two or three space dimensions, our modelling includes both surface tension and bending rigidity with volume conservation and allows the inclusion of external forces. We present a numerical method for the simulation of the model. Equations for and on the surface of the cell are discretised on a discrete surface. The efficacy of our methodology is illustrated by computer simulations of pseudopod driven migration and persistent migration in three space dimensions and the simulation of cell migration in the presence of obstacles.

While a major part of this work is the investigation of modelling generalisations through numerical simulation, especially with respect to cell motility in three-dimensions, our intention is to present a general modelling framework and numerical method that will be a potentially useful methodology for experimentalists and theoreticians alike in future studies of cell motility.

The remainder of our discussion proceeds as follows. In §2 we present our general modelling framework and our modelling assumptions. In §3 we present a numerical method for the approximation of surface evolution laws coupled to surface partial differential equations. In §4 we report the results of numerical simulations of a model for chemotaxis. In §5 we report the results of numerical simulations of a surface based model for the persistent motion of cells such as fish keratocytes. In §6 we discuss the implications of our findings in the study of cell motility and possible applications of our methodologies in future studies. We provide the technical details of our modelling and the numerical methods we employ in the electronic supplementary material (ESM).

2. A SURFACE BASED MODEL FOR CELL POLARISATION AND MOVEMENT

We consider models for cell motility and chemotaxis that consist of a geometric evolution law for a hypersurface representing the cell boundary coupled to a spatial pattern generator on the evolving surface describing polarisation of the cell. The particular form of the spatial patterning mechanism we shall investigate is a Turing pattern generator i.e., a semi-linear reaction diffusion system (RDS). The use of Turing type systems to model biological pattern formation phenomena is widespread (see [19] for a review) and recent numerical studies of Turing type systems on evolving surfaces show that while the key features of Turing mechanisms persist, such as spontaneous pattern formation and bifurcations due to surface evolution, the geometry of the evolving surface strongly influences the patterns expressed [20, 21]. We stress, that our general modelling strategy and the numerical methods we employ can be generalised to other possible polarisation mechanisms such as gradient based models or excitable network and wave based models all of which effectively couple surface partial differential equations to a surface evolution law [3].

2.1. Geometric evolution model. The cell membrane is represented by an evolving hypersurface, with the movement of the cell determined by the interaction of various forces that act normal to the cell membrane. We consider external forces such as a protrusive force associated with the RDS species and a pressure that constrains the enclosed volume, as well as internal forces that arise from the reaction of the cells surface to stretching and bending. We use the following force balance on the membrane, where we neglect the inertia of the membrane:

$$(2.1) \quad 0 = (\mathcal{F}_p + \mathcal{F}_v + \mathcal{F}_{visc} + \mathcal{F}_{ext} + \mathcal{F}_s + \mathcal{F}_b) \boldsymbol{\nu},$$

where $\boldsymbol{\nu}$ denotes the outward pointing unit normal to the surface Γ . We account for the following force contributions appearing in (2.1).

- A *protrusive force* depending on the densities of chemical species resident on the membrane (c.f. (2.9)) is denoted by

$$(2.2) \quad \mathcal{F}_p = \mathcal{F}_p(\mathbf{a}).$$

In the subsequent numerical simulations we make the phenomenological modelling assumption that the force is proportional to the species densities and given by $\mathcal{F}_p(\mathbf{a}) = \mathbf{k}_p \cdot \mathbf{a}$. The sign of the component $(k_p)_i$ of the vector \mathbf{k}_p governs whether the i 'th species promotes protrusion (positive) or retraction (negative) of the cell membrane. For $(k_p)_i > 0$ such a force may model the protrusive force generated by cross-linked filamentary actin in the vicinity of the cells surface while $(k_p)_i < 0$ could correspond to the contraction force generated by actin bundles [22].

- Experimental studies suggest that while the cell surface area may exhibit variability during movement the enclosed volume is relatively constant [18, 23]. We take this fact into account as a hard constraint, which means that the cell is able to immediately counterbalance small volume changes on the time scale of the RDS and the boundary evolution. In the following, a corresponding Lagrange multiplier will be denoted by $\lambda \in \mathbb{R}$. It can be interpreted as a pressure difference between interior and exterior of the cell. The corresponding force simply reads

$$\mathcal{F}_v = \lambda.$$

Note that the Lagrange multiplier λ is spatially constant and therefore models a spatially uniform force such that the enclosed volume is conserved.

- We include a *viscous force* that opposes motion

$$(2.3) \quad \mathcal{F}_{visc} = -\omega V,$$

where $\omega > 0$ is a kinetic coefficient and V is such that $V\boldsymbol{\nu} = \mathbf{V}$, where \mathbf{V} is the *material velocity* of the cell boundary (which we assume to be normal to the cell membrane). In the $2d$ case adhesion and de-adhesion may be modelled as an effective friction, i.e., a force, of the form (2.3), proportional to the local speed [24, 22]. In the $3d$ case the situation is more complicated and we intend to treat this subject in future work.

- We write \mathcal{F}_{ext} for any other *external normal forces* acting on the cell boundary where we have interaction with the medium in mind. As a concrete example and in order to illustrate the versatility of the proposed approach for cell motility in the complicated environments encountered in vitro, in §4.1.3 we present a simple phenomenological model for the movement of cells in the presence of obstacle particles that the cell cannot invade but that it may push out of its way.
- Resistance of the cell boundary to stretching may be incorporated by means of a *surface energy* of the form

$$(2.4) \quad \mathcal{E}_s = \int_{\Gamma} k_s$$

where $k_s > 0$ can be interpreted as a surface tension. The variation of the area functional is the mean curvature (we refer to [25] for details of a derivation), hence the force arising from the surface energy

is given by

$$(2.5) \quad \mathcal{F}_s = -k_s H,$$

where H is the mean curvature of Γ .

- The lipid bilayer forming the basic component of the cell membranes also resists bending. We consider the established model of [26] for the *bending energy*:

$$(2.6) \quad \mathcal{E}_b := \int_{\Gamma} \frac{k_b}{2} H^2,$$

where $k_b > 0$ is the bending rigidity. The variation of the bending energy yields the force contribution

$$(2.7) \quad \mathcal{F}_b = k_b \left(\Delta_{\Gamma} H + H |\nabla_{\Gamma} \boldsymbol{\nu}|^2 - \frac{1}{2} H^3 \right),$$

where ∇_{Γ} and Δ_{Γ} denote the surface gradient and Laplace-Beltrami operator respectively (ESM). We refer to [27] for a derivation.

Summing up the forces with their specific choices except for the external force we obtain the following equation for the evolution of the cell boundary:

$$(2.8) \quad \omega \mathbf{V} = \left(\mathbf{k}_p \cdot \mathbf{a} - k_s H + k_b \left(\Delta_{\Gamma} H + H |\nabla_{\Gamma} \boldsymbol{\nu}|^2 - \frac{1}{2} H^3 \right) + \lambda + \mathcal{F}_{ext} \right) \boldsymbol{\nu}.$$

The variational formulation of the evolution law (2.8), that we use to construct a finite element discretisation, is given in the ESM.

2.2. Cell polarisation model. We consider an RDS posed on an *evolving surface* $\{\Gamma(t)\}_{t>0}$:

$$(2.9) \quad \partial^{\bullet} \mathbf{a} + \mathbf{a} \nabla_{\Gamma(t)} \cdot \mathbf{V} - \mathbf{D} \Delta_{\Gamma(t)} \mathbf{a} = \mathbf{f}(\mathbf{a}) \quad \text{on } \Gamma(t), t > 0,$$

where $\mathbf{a} = (a_1, \dots, a_m)^t$, m is the number of chemical species involved, a_i denotes the density of the i th chemical species, \mathbf{V} is the material velocity of the surface (c.f., (2.3)),

$$(2.10) \quad \partial^{\bullet} \mathbf{a} := \partial_t \mathbf{a} + \mathbf{V} \cdot \nabla \mathbf{a},$$

is the material derivative with respect to the velocity \mathbf{V} , \mathbf{D} is a diagonal matrix of positive diffusion coefficients and $\mathbf{f}(\mathbf{a})$ is the reaction. For details of the derivation we refer for example to [10, 21]. We assume in the following that the evolving hypersurface is closed so that no boundary condition is required. For the initial condition we write

$$(2.11) \quad \mathbf{a}(\cdot, 0) = \mathbf{a}^0(\cdot) \quad \text{on } \Gamma(0).$$

3. DISCRETISATION

Here we describe the numerical methods we shall employ for the approximation of models for cell motility of the form described in §2. We keep the exposition relatively non-technical referring the interested reader to the ESM for the technical details. We decouple the approximation of the surface evolution and the RDS by treating the RDS concentration explicitly in the surface update step. The numerical method is based on approximating the surface $\Gamma(t)$ with a *triangulated surface* $\Gamma_h(t)$, which stems from the method described in the seminal paper of Dziuk [28].

3.1. Definition (Triangulated surface). A triangulated surface Γ_h is a polygon or polyhedron for $d = 2$ or 3 respectively with linear edges for $d = 2$ and planar faces for $d = 3$, such that

$$(3.1) \quad \Gamma_h = \cup_{s \in \mathcal{T}_h} s,$$

where \mathcal{T}_h consists of a finite number of closed intervals ($d = 2$) or a finite number of closed non-degenerate triangles ($d = 3$). For the simulations on surfaces ($d = 3$) we make use of *quadratic triangulated surfaces*. That is a surface that consists of curvilinear triangles each of which are the images of a reference triangle under a quadratic map as illustrated in Figure 1. We will use Γ_h to denote both




FIGURE 1. (Online version in colour.) Snapshots of the triangulation during a simulation of cell motility using a quadratic triangulated approximation to a surface.

a triangulated surface and a quadratic triangulated surface interchangeably. So as no confusion arises, we stress that for the approximation of smooth curves we consider triangulated (polygonal) discrete surfaces and for the approximation of smooth surfaces ($d = 3$) we consider quadratic triangulated surfaces. For details on triangulated surfaces, quadratic triangulated surfaces and approximation results we refer to [10, 29, 30, 31, 32].

The evolution law (2.8) is discretised using a surface finite element described in detail in the ESM. The method is based on the parametric finite element methods for fourth order geometric evolution equations derived in [33] and [11]. Under the proposed method the movement of the nodes of the triangulation satisfies the evolution law in the normal direction and includes a tangential velocity (that leaves the evolution law unchanged) which gives highly desirable mesh-properties in practice. Figure 2 illustrates an example of the robustness of the proposed scheme in the approximation of large deformations over schemes where nodes are moved solely in the normal direction.

To solve the RDS posed on the evolving surface we employ a surface finite element method based on the evolving surface finite element method [10] proposed by Dziuk and Elliot, where we account for the tangential velocity induced by the surface update scheme.

We also describe, in the ESM, a framework for the inclusion of stochastic external signals into the model. In particular for the subsequent simulations of chemotaxis we include a signal that is modelled by independent stochastic differential equations posed in each element (interval or triangle) which we approximate using the Euler-Maruyama method.

3.2. Software. The numerical methods were implemented using the adaptive finite element toolbox ALBERTA [34] and the linear systems were solved using UMFPACK [35], a direct solver for sparse linear systems.

4. MODELLING PSEUDOPOD DRIVEN CHEMOTAXIS

We investigate a model for eukaryotic chemotaxis originally proposed by Meinhardt [36]. The original model was posed at the discrete level and consisted of a three species RDS with a spatially varying local activator, a spatially varying local inhibitor and a spatially constant global inhibitor. Meinhardt proposed that such a model could account for the polarisation of chemotactic cells and the subsequent




FIGURE 2. (Online version in colour.) Snapshots of the mesh during two simulations of cell motility using a scheme that moves the nodes only in the normal direction and the scheme that includes a tangential velocity. The large deformations result in meshes that are not suitable for computation with the former approach necessitating a remeshing step while with the latter approach the mesh quality remains good throughout.

relocation and splitting of activator peaks in response to changing external signals. He did not however consider the mechanical aspects of the evolution of the cell membrane. Neilson et al. [7] investigated a continuous form of the model where the three species were all spatially dependent, approximating the model equations with a surface finite element method for the RDS approximation and a level set method for the surface update. They have conducted detailed comparisons of their simulations using a level set method with experiments [37] as well as some preliminary investigations into robust computational methods, specifically short time simulations using a surface finite element method [16]. All their modelling and simulation was conducted in $2d$, the model we consider extends the previous work by increasing the dimension, accounting for the bending energy and modelling obstacles.

In the original model posed by Meinhardt, the spatial independence of the global inhibitor is used in the derivation of the model. Since the global inhibitor is spatially constant, its concentration can be obtained by averaging, i.e., the use of a non-local term (the mean value of the local activator). We therefore consider the following transformation of Meinhardt's model from the spatially discrete fixed surface setting, to a continuous evolving surface:

$$\begin{aligned}
 (4.1) \quad & \partial^\bullet a_1 + a_1 \nabla_{\Gamma(t)} \cdot \mathbf{V} - D_1 \Delta_{\Gamma(t)} a_1 = \gamma \left(\frac{(r_1 + s)(a_1^2/a_2 + b_1)}{(s_3 + a_3)(1 + s_1 a_1^2)} - r_1 a_1 \right), \\
 & a_2 = \frac{1}{|\Gamma(t)|} \int_{\Gamma(t)} a_1, \\
 & \partial^\bullet a_3 + a_3 \nabla_{\Gamma(t)} \cdot \mathbf{V} - D_3 \Delta_{\Gamma(t)} a_3 = \gamma(b_3 a_1 - r_3 a_3), \quad \text{on } \Gamma(t), t > 0 \\
 & \mathbf{a}(\cdot, 0) = \mathbf{a}^0(\cdot) \quad \text{on } \Gamma_0.
 \end{aligned}$$

Here the r_i, s_i and b_i 's are material parameters, γ is a scaling parameter that governs the overall timescale of the reaction rate and the term $s(\mathbf{x}, t)$ models both the underlying noise from the external media and a noisy chemotactic signal. The term involving the chemotactic signal feeds multiplicatively into the autocatalytic term in the activator equation. Weak signals are amplified due to autocatalysis, thus the model provides a mechanism for gradient sensing of small chemotactic gradients. The RDS is coupled to the surface evolution law solely through the activator concentration a_1 which promotes protrusion of

Reaction kinetic parameters								
D_1	D_3	γ	r_1	r_3	s_1	s_3	b_1	b_3
1.0	7.0	2.5×10^4	2×10^{-2}	13×10^{-3}	1×10^{-4}	0.2	0.1	5×10^{-3}

TABLE 1. Parameter values for numerical experiments of the movement of two-dimensional cells with the Meinhardt kinetics 4.1.

the cell membrane. Adopting the notation of §2, the contribution of the RDS species to the evolution law (c.f. (2.2)) reads

$$(4.2) \quad \mathcal{F}_p = \mathbf{k}_p \cdot \mathbf{a} = (k_p)_1 a_1.$$

In fact the activator and inhibitor are in phase and this is one potential drawback of the Meinhardt model as it presents no obvious mechanism for coupling surface concentrations to retraction of the cell membrane. One can show that the model (4.1) is equivalent to the model considered by Neilson et al., if the diffusivity of the global inhibitor in the Neilson model is sufficiently large relative to the diffusivity of the other two species [38].

We model the stochastic term $s(\mathbf{x}, t)$ in (4.1) as the sum of an underlying noise term $\eta^t(\mathbf{x})$ due to, say, heterogeneity in the medium and a term $R^t(\mathbf{x})$ that models the cells sensing of the chemotactic signal, i.e., $s(\mathbf{x}, t) := (R^t(\mathbf{x}) + \eta^t(\mathbf{x}))$. The underlying noise term (present in all the simulations) satisfies the following Ornstein-Uhlenbeck (mean-reversion) stochastic process

$$(4.3) \quad d\eta^t = -\eta^t dt + 2 \times 10^{-4} dW^t,$$

where W^t denotes the Wiener process. We discretise in space by assuming η is constant over each finite element and the Euler Maruyama method is used to approximate the solution numerically (for details see the ESM).

4.1. Results. For the results on curves we took the unit circle as the initial steady state and used the reaction kinetic parameter values given in Table 1. The parameter values are those Meinhardt used in his original study rescaled such that the diffusivity of the activator is 1.0, the only parameter we have tuned is the saturation of the activator concentration s_1 , which is smaller to account for dilution in the activator concentration due to domain growth. For all the simulations on curves we used the same equidistributed initial mesh with 1024 degrees of freedom (further refinement did not change the solutions qualitatively) and a timestep of 10^{-5} . The CPU times of all the calculations on curves is in the order of minutes, for example the two simulations reported in Figure 3 had CPU times of just over 2000 seconds.

4.1.1. Random migration. Figure 3 shows the centroid trajectories of 5 cells migrating under two different geometric evolutions, surface tension evolution ($k_s = 25, k_b = 0, (k_p)_1 = 1.5$) and combined surface tension - elastic evolution ($k_s = 23, k_b = 3, (k_p)_1 = 2$), with no chemotactic signal and different realisations of the noise term. In both cases, the cells initially polarise and then migrate, with roughly two pseudopods at the front of the cell at any given time. The cells change direction via splitting and decay of pseudopods with one pseudopod splitting and persisting while the other decays. This leads to the characteristic linear motion over short times interspersed with sharp changes in direction corresponding to a pseudopod splitting/decay event, similar to that observed by Neilson et al. [7]. Under surface tension evolution the cells maintain a characteristic shape (two pseudopods at the front of the cell with a valley between them), while the introduction of bending rigidity generates a greater variety of cell shapes and the cell no longer has a characteristic shape with varied banana like shapes evident (the cell shapes resemble those in the simulations presented in Figure 5). From the simulations we observe that the diluting effect of protrusion plays an important role in destabilising activator peaks and pseudopod splitting. By this we refer to the fact that the formation of an activator peak results in protrusion of the cell membrane which in turn leads locally to an decrease in activator density (as protrusion may be viewed as volumetric

Parameter	Value	Physical interpretation
r_0	1	Initial radius of cell sets a length scale of $4\mu m$.
$(k_p)_1$	$1.5 - 2$	Coefficient of protrusive forcing term implies a timescale of $\sim 1800s$.
D_1	1.0	Implies a diffusivity of $2.22 \times 10^{-3} \mu m s^{-1}$.
D_3	7.0	Implies a diffusivity of $1.55 \times 10^{-2} \mu m s^{-1}$.
k_s	~ 25	Assuming a surface tension of $1pN$ this sets a kinetic scale for the simulations.
k_b	3	Implies a bending rigidity of approximately $1.92pN \mu m^2$.

TABLE 2. Physically relevant parameter values for simulations on curves.

expansion). As the local maxima corresponding to the activator peak is reduced most at the tip of the peak, where protrusion is largest, this has the effect of increasing the propensity of activator peaks and hence pseudopods to split.

We may proceed to estimate some of the parameter values using available experimental data, which is readily available for dictyostelium cells [2]. The typical radius of cell cross sections is $4\mu m$ which sets the length scale for the computations. The maximum actin polymerisation velocity (which is related to the nondimensional parameter $(k_p)_1$) is approximately $0.1ms^{-1}$, thus the value of $(k_p)_1$ together with the maximum density of a_1 in the simulations (approx. 30) sets the timescale for the simulations. Typical values of the surface tension are $10pN/\mu m$, assuming a cell height of $0.1\mu m$ this sets a kinetic scale for the simulation. The remaining physically relevant parameters may thus be estimated and are given in Table 2. Note that the timescale we refer to in Table 2 corresponds to one unit of computational time. The length of the simulations in Figure 3 is one computational time unit or thirty minutes in actual time and corresponds to roughly 20 pseudopod lifetimes (Each change in direction in Figure 3 represents a pseudopod splitting/decay event), thus the timescale of an individual pseudopod is around ninety seconds.

We note that other choices of the material parameters, specifically weaker surface tension, gives cells with more elongated shapes, larger protrusions and cell bodies which appear less rounded.

4.1.2. Migration in the presence of a chemoattractant. We now include a chemoattractant in the model. We use the stochastic receptor model proposed by Neilson et al. [7] to model the noisy chemotactic signalling. For completeness we state the essential details. At time $t \in (0, T]$ they model the cells sensing of the chemotactic signal R^t with an Ornstein-Uhlenbeck stochastic process of the form

$$(4.4) \quad dR^t = \theta(\mu - R^t) dt + \sigma dW^t,$$

where W^t denotes the Wiener process, $\mu(\mathbf{x}, t)$ models the strength of the chemotactic signal, $\theta(\mathbf{x}, t)$ the rate of reversion to the mean μ and $\sigma(\mathbf{x}, t)$ the variance. The mean μ is local and prescribed by the model, while the rate of reversion to the mean and variance are local too as they are chosen such that $\theta = 1/(1 - \mu)$ and $\sigma = c\mu^{1/2}$ (for details see [7, §6.2.2]). To compare with [7, §6.2.5], we model the signal such that if a chemoattractant is present the mean μ varies from the base signal strength (at the back of the cell) of 0.5 to $0.5 + \rho$ at the front, where $\rho > 0$ represents the signal strength. For a given signal direction \mathbf{d}_s the position of the rear of the cell \mathbf{x}_r is such that $\mathbf{x}_r \cdot \mathbf{d}_s = \min_{\Gamma}(\mathbf{x} \cdot \mathbf{d}_s)$. We have also conducted experiments where $\mu = \exp(-c|\mathbf{x} - \mathbf{x}_c|)$, where \mathbf{x}_c denotes the location of a static point source of chemoattractant and observe similar results. We discretise (in space) by assuming the mean μ is constant over each finite element, and hence θ and σ are also constant on each finite element.

Figure 4 shows the trajectories of the centroids of 5 cells migrating leftwards in a linear chemotactic gradient of varying strength under conserved surface tension evolution ($k_s = 25, k_b = 0, (k_p)_1 = 1.5$). The results are similar under the other geometric evolution considered ($k_s = 22, k_b = 3, (k_p)_1 = 2$) and are not illustrated. The migration of the cells with only the base signal and the signal strength set to zero is reported in Figure 5(a). We observe no clear directional preference similar to the migration observed in




FIGURE 3. (Online version in colour.) Centroid trajectories of 5 cells migrating in the absence of a chemoattractant under different geometric evolution laws. In both cases we see motion in a straight line for short times punctuated by sharp changes in direction corresponding to pseudopod splitting/decay.

the absence of a chemoattractant. As the signal strength is increased (at a signal strength of 0.04 around 8% of the base signal) the cells start to exhibit a clear directional preference and successfully navigate up the chemotactic gradient. In Table 3 we report on chemotaxis measures of 100 cells migrating under the 6 different signal strengths shown in Figure 4. We state the average value over the 100 simulations (for each signal strength) of the following quantities all evaluated at $t = 0.5$:

- The chemotactic index (CI), defined as the cosine of the angle between a line connecting the present position of the cell centroid to the starting point and a line directly up the chemotactic gradient [39].
- The persistence length (PL) of the centroid trajectory in the x and y directions. The persistence length is taken to be the displacement in the chosen direction divided by the total length of the trajectory of the cell centroid [40].
- The squared displacement of the cell centroid from its initial position.
- The speed of the cell.

The data suggests that as the strength of the chemotactic signal is increased the cells exhibit greater propensity for persistent migration up the chemical gradient with chemotactic indices similar to those observed experimentally in the case of *Dictyostelium* cells [39, 0.71-0.94]. The persistence length in the x direction (up the chemotactic gradient) also increases with the signal strength while the persistence length in the y direction is reduced. We also note increasing the signal strength leads to larger displacements. The results suggest that for values of $\rho \geq 0.06$ the cell is able to migrate successfully up the chemical gradient with all the reported statistics converging to similar means for further increases in the signal

Signal strength	CI	PL (x)	PL (y)
0	N/A	0.4336 (0.2346)	0.4601 (0.2442)
0.02	0.7196 (0.2877)	0.4938 (0.2188)	0.3418 (0.1999)
0.04	0.9423 (0.0742)	0.6968 (0.1177)	0.2005 (0.1163)
0.06	0.9888 (0.0133)	0.8510 (0.0511)	0.1088 (0.0685)
0.08	0.9860 (0.0120)	0.8490 (0.0350)	0.1288 (0.0676)
0.1	0.9898 (0.0141)	0.8489 (0.0272)	0.0987 (0.0734)
Signal strength	Squared displacement (arb)	Speed (arb)	Speed ($\mu\text{m/s}$)
0	37.052 (14.5937)	20.056 (1.9781)	0.0445
0.02	34.430 (13.5045)	19.799 (2.8624)	0.0440
0.04	44.655 (11.9692)	20.077 (2.8633)	0.0446
0.06	66.011 (10.3248)	21.156 (1.9327)	0.0470
0.08	66.491 (6.8996)	21.524 (1.9714)	0.0478
0.1	67.387 (8.8005)	21.435 (2.1560)	0.0476

TABLE 3. Mean and standard deviation (in parentheses) of chemotaxis measures at $t = 0.5$ for 100 cells migrating as in Figure 4.

strength. The (physical) cell speeds are similar to those observed in migrating leukocytes [41, Tab. 1] and Dictyostelium cells [39] (in both cases reported in $\mu\text{m}/\text{min}$).

We now investigate the ability of this model to capture the ability of a cell to respond to a changing chemotactic signal. We use the same stochastic receptor model for the chemotactic signalling but now we change the direction of the signal at various stages in the evolution. Figures 5(a) and 5(b) show snapshots of the cells shaded by activator concentration under the two different geometric evolutions ($k_s = 25, k_b = 0, (k_p)_1 = 1.5$ and $k_s = 22, k_b = 3, (k_p)_1 = 2$) in response to a changing chemotactic signal. Initially we include only the base signal with noise, i.e., the signal strength is set to zero. At the times in the evolution at which the arrows appear in the figure we change the direction of the signal, with signal strength $\rho = 0.1$, to the direction indicated by the arrows. We see that under both geometric evolutions the cell successfully responds to the changing signal exhibiting a clear directional preference for movement in the direction of higher chemoattractant concentration. As a final example of response to a changing signal, we consider the case where the signal direction is changed by 180 degrees. The results of such a simulation are shown in Figure 6. We observe the cell successfully responds to the change in signal direction and does so via turning gradually through 180 degrees. This corresponds to so called “hops” (consecutive right/right or left/left splitting of pseudopods) that are an important mechanism for the reorientation of Dictyostelium cells moving in a direction more than 90 degrees off the chemotactic gradient [42, Fig. 4]. Under this model, we have however thusfar not observed the formation of *de novo* pseudopods towards the direction of increasing chemoattractant, which are another significant mechanism for major directional corrections [42].

4.1.3. Migration in the presence of obstacles. We now include an external force in the evolution law which arises from a model for the migration of cells through a field of obstacles. We model the obstacle particles as rigid spherical bodies. The obstacle-cell interaction is described by a phenomenological repulsive force that points in the direction *normal* to the cell membrane with no tangential component. Unlike the models proposed by Hecht et al. [43] and Grima [44] the obstacles and the cell both move due to mechanical interactions.




FIGURE 4. (Online version in colour.) Centroid trajectories of 5 cells migrating leftwards in the presence of a linear chemoattractant gradient under conserved surface tension evolution with varying signal strength (ρ).

Let $N_o \in \mathbb{N}$ denote the number of obstacles with centres $\{\mathbf{m}_i\}_{i=1}^{N_o}$ and radii $\{r_i\}_{i=1}^{N_o}$. For the force acting on a point $\mathbf{x} \in \Gamma(t)$ on the cell boundary due to the interaction with obstacle i we postulate

$$(4.5) \quad \mathcal{F}_{o,i} = \max(0, (r_i(1 + \epsilon) - |\mathbf{m}_i - \mathbf{x}|)) ((\mathbf{m}_i - \mathbf{x}) \cdot (-\boldsymbol{\nu})) \frac{f_i}{|\mathbf{m}_i - \mathbf{x}| - r_i}$$

where $f_i > 0$ is a material coefficient and $\epsilon > 0$ is a thickness parameter: the force is zero if the distance between the cell membrane and the obstacle boundary is bigger than ϵr_i . The force becomes infinite as this distance approaches zero and then dominates any other forces on the cell membrane, thus preventing intersection of the cell and the obstacle. The external force acting on the cell boundary is given by

$$(4.6) \quad \mathcal{F}_{ext} = \sum_{i=1}^{N_o} \mathcal{F}_{o,i}$$

For the obstacle particles we postulate a viscous motion law, too, where the reaction forces from the cell boundary $-\mathcal{F}_{i,o}$ and obstacle-obstacle interactions are taken into account. We postulate

$$(4.7) \quad \omega_i \dot{\mathbf{m}}_i = - \int_{\Gamma(t)} \mathcal{F}_{i,o} \boldsymbol{\nu} dS + \sum_{j \neq i} \mathcal{F}_{j,i}.$$

Here, the $\omega_i > 0$ are positive kinetic coefficients related to the mass of the particle, the first term on the right hand side modelling the cell-obstacle interaction is

$$(4.8) \quad \int_{\Gamma(t)} \mathcal{F}_{i,o} \boldsymbol{\nu} dS = \int_{\Gamma} \max(0, (r_i(1 + \epsilon) - |\mathbf{m}_i - \mathbf{x}|)) ((\mathbf{m}_i - \mathbf{x}) \cdot \boldsymbol{\nu}) \frac{f_i}{|\mathbf{m}_i - \mathbf{x}| - r_i} \boldsymbol{\nu} dS,$$




FIGURE 5. (Online version in colour.) Response to a changing chemotactic signal. Initially there is no signal with arrows indicating the time at which a signal is introduced and the signal direction. Note the two figures are not on the same scale and the cells have the same enclosed volume.




FIGURE 6. (Online version in colour.) Response to a changing chemotactic signal. In this example the direction of the chemotactic signal is changed by 180 degrees. We observe that the cell turns through 180 degrees and successfully responds to the changing signal, migrating up the new chemotactic gradient. Cell outlines shown every 0.1 units of computational time over the interval $[0, 1]$.

Reaction kinetic parameters								
D_1	D_3	γ	r_1	r_3	s_1	s_3	b_1	b_3
10	70	5×10^4	2×10^{-2}	13×10^{-3}	1×10^{-4}	0.2	0.1	5×10^{-3}

TABLE 4. Parameter values for numerical experiments of the movement of three-dimensional cells with the Meinhardt kinetics 4.1.

Parameter	Value	Physical interpretation
r_0	1	Initial radius of cell sets a length scale of $1.17\mu m$.
$(k_p)_1$	0.5	Coefficient of protrusive forcing term implies a timescale of $\sim 230s$.
D_1	10	Implies a diffusivity of $5.95 \times 10^{-2} \mu m^2 s^{-1}$.
D_2	70	Implies a diffusivity of $4.17 \times 10^{-1} \mu m^2 s^{-1}$.
k_s	~ 25	Assuming a surface tension of $10pN/\mu m$, sets kinetic scale.

TABLE 5. Physically relevant parameter values for simulation of migration of a cell in three dimensions.

and the $\mathcal{F}_{j,i}$ is the force from particle j exerted on particle i for which we postulate

$$(4.9) \quad \mathcal{F}_{j,i} = \max(0, (1 + \epsilon)(r_i + r_j) - |\mathbf{m}_i - \mathbf{m}_j|) \frac{f_{ji}}{|\mathbf{m}_i - \mathbf{m}_j| - (r_i + r_j)} (\mathbf{m}_i - \mathbf{m}_j)$$

where the $f_{ji} = f_{ij} > 0$ are material coefficients. Note that in the absence of the cell, the initial location of the obstacles is such that the sum of the forces $\mathcal{F}_{j,i}$ yields zero so that the particles do not move. Moreover we have the following balance of forces exerted by the cell on the obstacles and forces on the cell membrane due to the obstacles

$$\int_{\Gamma} (\mathcal{F}_{ext}\nu) + \sum_{i=1}^{N_o} \left(\int_{\Gamma} (-\mathcal{F}_{i,o}\nu) + \sum_{j \neq i} \mathcal{F}_{j,i} \right) = 0.$$

Figures 7 and 8 show a series of snapshots of cell migration through a field of obstacles, with parameter values as in Table 1 and the two previously considered geometric evolutions. Our numerical experience suggests that under the simple model of cell obstacle interactions we have employed, the increase in computational time, even with a large number of obstacles, from the case of no obstacles is negligible. We include the forcing terms in the evolution law for the cell membrane and the obstacle centres given by (4.6)–(4.9), with parameter values $\epsilon = 0.1$, $f_i = f_{ij} = 100$ for all i, j and $\omega_i = r_i/100$. We observe that the cell successfully migrates through the field of obstacles maintaining the characteristic shape as it deflects the obstacles. Our numerical experiments suggest that this behaviour is sensitive to the parameter values chosen in the repulsive potential (4.5). In particular, if we set the kinetic coefficient related to the mass of the obstacles ω_i c.f., (4.8) to be comparable in magnitude to the kinetic coefficient related to the mass of the cell (1 by assumption), which means the obstacles inhibit more strongly the protrusion of pseudopods, then pseudopod splitting no longer occurs and the cell exhibits persistent motion in the direction of an obstacle (not reported).

4.1.4. Migration of cells in three space dimensions. We now present results for the motion of three-dimensional cells in the absence of a chemoattractant. We took the unit sphere as the initial steady state and used the reaction kinetic parameter values given in Table 4. We selected a timestep of 10^{-5} and used the adaptive strategy described in the ESM with parameters $N_H = 0.5$, $N_h = 0.75$, $M_H = 0.25$ and $M_h = 0.5$. We considered an evolution law of the form (2.8) with parameters $k_s = 25$, $k_b = 0$, $(k_p)_1 = 0.5$. As in Table 2, we give a physical interpretation of the parameter values in Table 5, assuming the



FIGURE 7. (Online version in colour.) Undirected migration (i.e., migration in the absence of a chemoattractant) in the presence of obstacles of a cell under conserved surface tension evolution, for parameter values see Table 1 and text.

radius of the spherical cell at rest is $1.17\mu m$. The model for the random signalling was the same as the model used in the two-dimensional case. Snapshots of the cell surface shaded by activator concentration are reported in Figure 9. We see qualitatively similar behaviour to the case of curves with protrusion of activator peaks leading to pseudopod formation and the rest of the cell retracting behind the pseudopods.




FIGURE 8. (Online version in colour.) Undirected migration in the presence of obstacles of a cell under a combination of conserved surface tension and elastic evolution, for parameter values see Table 1 and text.

We also observe pseudopod splitting as the cell changes direction via biased generation and retraction of existing pseudopods. The simulation is considerably more challenging than the curve case considered previously and the total CPU time of the simulation was just over 27 hours.




FIGURE 9. (Online version in colour.) Migration in the absence of a chemoattractant of a three-dimensional cell under conserved surface tension evolution, for parameter values see Table 4 and text.

5. MODELLING THE PERSISTENT MOTION OF KERATOCYTES

We present a model based on the general modelling framework described in §2 that seeks to capture the persistent motion of fish keratocytes. The cells deform rapidly into a temporally persistent shape and once in this shape move at a constant speed without changes in direction. Keren et al. [18] conduct an analysis of the shapes taken by moving keratocyte cells and propose a simple phenomenological model to account for the observed movement and cell shapes. Their results suggest that the steady state shapes of the cell are broadly described by two modes and that cell shape, specifically the aspect ratio (length/width), is strongly correlated with the speed of motion. They also examined the actin distribution within the cell. Branched actin filaments promoting protrusion are concentrated at the fast moving front and retraction promoting actin bundles are concentrated at the rear. The steady state appears stable to perturbations and

Reaction kinetic parameters				Surface evolution parameters			
D_1	D_2	γ	k_1	k_s	k_b	$(k_p)_1$	$(k_p)_2$
0.5	50	10	0.1	2	2	-2	1

TABLE 6. Parameter values for numerical experiments of keratocyte movement with the RDS (5.1).

if movement of the cell is disrupted the cell rapidly regains its previous shape and speed of movement, usually moving in a new direction.

The observed behaviour of spontaneous polarisation and subsequent development of a steady state stable to perturbations, suggests a Turing type mechanism coupled to a surface evolution law could accurately capture the observed dynamics. Shao et al. [22] considered a membrane subject to surface tension, bending rigidity, and forcing with volume conservation. The forcing strength was dependent on the concentrations of a two component RDS posed in the bulk of the cell. They present computational results, for two-dimensional cells with weak volume conservation (enforced via penalisation), based on a phase field method. Ziebert et al. [6] present a model for keratocyte movement, again based on a phase field method, where they couple the surface evolution to a vector field that seeks to describe the polarisation of the actin network. Studies suggest that branched filamentary actin and actin bundles are concentrated primarily near the cell membrane near areas of protrusion and retraction respectively, while away from the cell surface the actin is in a remodelling phase between that of branched and bundled actin [18, 45, 46]. This suggests a surface model where the pattern formation process occurs on the cell membrane itself may be appropriate. We propose the *activator-depleted* substrate model [47]:

$$\begin{aligned}
 (5.1) \quad & \partial^\bullet a_1 + a_1 \nabla_{\Gamma(t)} \cdot \mathbf{V} - D_1 \Delta_\Gamma a_1 = \gamma (k_1 - a_1 + a_1^2 a_2), \\
 & \partial^\bullet a_2 + a_2 \nabla_{\Gamma(t)} \cdot \mathbf{V} - D_2 \Delta_\Gamma a_2 = \gamma (k_2 - a_1^2 a_2), \quad \text{on } \Gamma(t), t > 0, \\
 & \mathbf{a}(\cdot, 0) = \mathbf{a}^0(\cdot) \quad \text{on } \Gamma_0.
 \end{aligned}$$

We first present results for curves, with material and RDS parameters given in Table 6. We considered an initially circular cell with radius 1 centred at the origin. The initial condition for the RDS was taken as the linearly stable steady state $a_1^0 = k_1 + k_2$, $a_2^0 = k_2 / (a_1^0)^2$ with a symmetry breaking perturbation of the form $\max(1 \times 10^{-4} x_1, 0)$ added to the initial condition of the a_2 species. The specific form of the initial condition leads to cells that migrate only along the x axis (we verified that the choice of other initial conditions only changed the direction of the movement). The hypothesis of Keren et al. is that variability in the actin dynamics is the major factor governing the observed variations in shape and speed. To investigate this hypothesis, we propose that the a_1 species in the RDS (5.1) corresponds to the density of retraction promoting actin bundles while the a_2 species corresponds to the density of protrusion promoting actin filaments, which is similar to the model considered in [22]. We can model variable actin dynamics by changing the constant k_2 which can be interpreted as the growth rate of actin filaments. Increasing k_2 leads to higher concentrations of a_2 relative to a_1 and thus should lead to faster moving cells with stronger forcing at the front.

In all the simulations on curves we used an initially equidistributed mesh with 1024 degrees of freedom and a fixed timestep of 10^{-3} . The CPU times were on the order of seconds with a typical simulation taking approximately 200 seconds. Figures 10(a) and 10(b) show, for different values of k_2 , the initial position of the cells at time 0 and the cell positions and surface RDS concentrations at time 5 (by which time all the cells have reached a steady state with constant speed and time independent RDS concentrations). We see faster cell speeds and larger aspect ratios for increased values of k_2 , similar to the models where the RDS is posed in the bulk [22]. The shapes of the cells at steady state also resemble those observed experimentally [18]. In particular we see the rounded “D” shape in the right most cell corresponding to $k_2 = 0.6$ and the much more elongated “canoe” shape in the left most cell corresponding to $k_2 = 1.8$.

Parameter	Value	Physical interpretation
r_0	1	Initial radius of cell sets a length scale of $4\mu m$.
$(k_p)_2$	1	Coefficient of protrusive forcing term implies a timescale of $\sim 45s$.
D_1	0.5	Implies the diffusivity of actin bundles of $4.44 \times 10^{-2} \mu m s^{-1}$.
D_2	50	Implies the diffusivity of actin filaments of $4.44 \mu m s^{-1}$.
k_2	.6 – 1.8	Implies an actin filament growth rate of $1.33 \times 10^{-2} - 4 \times 10^{-2} s^{-1}$.
k_s	2	Assuming a surface tension of $1pN$ sets the kinetic scale.
k_b	2	Implies a bending rigidity of approximately $16pN \mu m^2$.

TABLE 7. Physically relevant parameter values for simulation of keratocyte movement in two-dimensions.




FIGURE 10. (Online version in colour.) Initial position (at $t = 0$ right hand cell) and persistent keratocyte like migration of cells (at $t = 5$). The parameter $k_2 = 0.6, 0.8, 1.0, 1.2, 1.4, 1.6, 1.8$ reading from right to left for the 7 polarised (left hand) cells (c.f. (5.1)) with the remaining parameters given in Table 6.

We report on the aspect ratio $AR = (\lambda_2/\lambda_1)^{1/2}$, as considered in [6], where for $i = 1, 2$, λ_i is as follows (the λ_i 's are the eigenvalues of the diagonal 2×2 variance matrix of the cells centroid):

$$\lambda_i = \frac{1}{3} \int_{\Gamma} (x_i - x_i^c)^2 \nu_i ds,$$

where x_i^c is the i th coordinate of the cells centroid. We also report on the deviation from reflection symmetry of the migrating cells as considered in [6]. This is measured by the following quantities (the nonzero components of the skewness tensor of the cells centroid scaled by a constant factor):

$$\eta_1 = \left(\frac{1}{4} \int_{\Gamma} (x_1 - x_1^c)^4 \nu_1 ds \right)^{1/3} (\lambda_1 + \lambda_2)^{-1/2},$$

$$\eta_2 = \left(\frac{1}{2} \int_{\Gamma} (x_1 - x_1^c)^2 (x_2 - x_2^c)^2 \nu_1 ds \right)^{1/3} (\lambda_1 + \lambda_2)^{-1/2}.$$

Figure 11 shows plots of the speed of the cell centroids and the aspect ratio of the cells both against time. We clearly see the positive relationship between aspect ratio and cell speed evident in the experimental




FIGURE 11. (Online version in colour.) The speed of the cell centroid and aspect ratio both vs. time of the cells shown in Figure 10. We observe a positive relationship between aspect ratio and speed.




FIGURE 12. (Online version in colour.) Asymmetry measures vs. time of the cells shown in Figure 10. We observe larger deviations from reflection symmetry in the cells travelling at slower speeds. As the cells attain persistent shapes the values converge to a steady state.

studies. In physical units the range of the speed at steady state of the cells shown in Figure 10 is 0.178 to $0.445 \mu\text{m s}^{-1}$. Both the speed and aspect ratios are similar to those observed in the experimental results reported in [18, Figure 4b]. Figure 12 shows values of the asymmetry measures against time. We see that the cells travelling at slower speed exhibit larger deviations from reflection symmetry. This is in contrast to the results obtained under the model considered in [6, Fig. 4], where η_1 does increase as the speed of the cells decreases but η_2 is positively correlated with speed. We note that for the first two simulations $k_2 = 0.6$ and 0.8 , η_2 is negatively correlated with speed. As an ellipse satisfies $\eta_1 = \eta_2 = 0$, it is not clear that the faster moving cells with larger aspect ratios, but more elliptical profiles, should exhibit larger deviations from reflection symmetry. We also observe that after a brief initial stage in which all the

reported values oscillate, the values converge to a steady state as the cells travel in a persistent fashion with a fixed shape and a constant speed.




FIGURE 13. (Online version in colour.) Initial position (at $t = 0$ right hand cell) and persistent keratocyte like migration of cells (at $t = 5$). The parameter $k_2 = 0.6, 0.7, 0.8, 0.9, 1.0$ reading from right to left for the 5 polarised (left hand) cells (c.f. (5.1)), for the remaining parameter values see Table 8.




FIGURE 14. (Online version in colour.) The speed of the cell centroid and cell surface area both vs. time of the cells shown in Figure 13. We observe a positive relationship between surface area and speed.

We also report on simulations of three-dimensional keratocyte motion. We took the unit sphere as the initial cell shape, the same initial conditions for the RDS concentrations as in the curve case, a fixed

Reaction kinetic parameters				Surface evolution parameters				Adaptive strategy parameters			
D_1	D_2	γ	k_1	k_s	k_b	$(k_p)_1$	$(k_p)_2$	N_H	N_h	M_H	M_h
1	100	20	0.1	1	1	-0.7	.35	.5	1	.25	.5

TABLE 8. Parameter values for numerical experiments of three-dimensional keratocyte movement with the RDS (5.1) and surface evolution law 2.8.

Parameter	Value	Physical interpretation
r_0	1	Initial radius of cell sets a length scale of $1.17\mu m$.
$(k_p)_2$.35	Coefficient of protrusive forcing term implies a timescale of $\sim 4s$.
D_1	1.0	Implies the diffusivity of actin bundles of $0.342\mu m^2 s^{-1}$.
D_2	100	Implies the diffusivity of actin filaments of $34.2\mu m^2 s^{-1}$.
k_2	.6 – 1.0	Implies an actin filament growth rate of $0.15 \times 10^{-1} - 0.25s^{-1}$.
k_s	1	Assuming a surface tension of $10pN\mu m^{-1}$, sets the kinetic scale.
k_b	1	Implies a bending rigidity of approximately $13.69pN\mu m$.

TABLE 9. Physically relevant parameter values for simulation of keratocyte movement in three-dimensions.

timestep of 10^{-4} and remaining parameter values for both the surface evolution and adaptive strategy as given in Table 8. The CPU times were on the order of minutes with a typical simulation taking approximately 2000 seconds. Proceeding as in §4, we give physical interpretations of the parameter values for curves and surfaces in Tables 7 and 9 respectively. Figures 13(a) and 13(b) show a similar experiment to the one carried out for curves now on surfaces, specifically we report for different values of k_2 , the initial position of the cells at time 0 and the cell positions and surface RDS concentrations at time 5 (by which time all the cells have reached a steady state with constant speed and time independent RDS concentrations). The gross behaviour is the same as the curve case, in that as the parameter k_2 is increased from 0.6 to 1 the cells move faster at steady state and appear more elongated. Figure 14 shows plots of the speed of the cell centroids and the surface area of the cells both against time. We plot the surface area as it is proportional to the two, roughly equal, aspect ratios in the (x, y) and (x, z) directions. We observe the same positive relationship as in the curve case with both surface area and speed converging to steady states. We have also verified that the aspect ratios converge to steady states with the aspect ratio in the (y, z) direction approaching 1. Note for larger values of k_2 the cells developed a self intersection which is inadmissible under our modelling as it would correspond to a change in topology in the physical setting. Scenarios where one wishes to consider topological change, or respectively methods that avoid topological change, are a subject of our current research.

6. CONCLUSION

In this work we have presented a computational framework for the modelling of cell motility. We propose a simple and consistent means of coupling cell movement with gradient sensing, polarisation using surface PDEs and external forces. Our methods can be generalised to the modelling of more complex phenomena such as adhesion and crawling on a substrate or cell-cell interactions and we illustrate one such generalisation with a concrete example of migration in the presence of obstacles.

A contribution of our study is the description of a numerical method for the simulation of cell movement that can account for the large deformations that arise in simulations of cell motility and that can be applied to the study of three-dimensional cell migration. The model equations consist of PDEs for and on surfaces and the numerical method seeks to approximate these equations on a discrete surface. Thus

both the continuous and discrete problems are posed in one dimension less than the underlying spatial dimension, which in the case of the discrete problem typically means fewer degrees of freedom are needed than would be the case for embedded methods [10, 13, 14, 15, 16]. On curves, our experience is that the method maintains a mesh suitable for computation without the use of remeshing or adaptive mesh refinement. For the simulations of cell migration in three-dimensions we occasionally observed deterioration in the mesh quality, even with the redistribution of the vertices implicit in our numerical scheme, necessitating spatial adaptivity. An area of our ongoing research is the investigation of numerical methods robust to large deformations in the cell surface.

We consider a pseudopod centred model for chemotaxis similar in form to that considered in [37]. Unlike compass models [39], which are reasonable for cells with flexible polarity where a large gradient may induce pseudopods at any position on the cell membrane, pseudopod centred models [17] are suitable for strongly polarised cells where pseudopods are generated preferentially at the front with directional bias, due to a chemotactic gradient, restricted primarily to small changes in direction [42]. The major contributions and imports of our study are the inclusion of bending rigidity, the inclusion of external forces, the observation that the gross behaviour, of pseudopod splitting, observed in [37] for $2d$ cells persists in $3d$ simulations and that the model remains qualitatively unchanged when one considers a two component RDS, with a spatially constant global inhibitor, rather than a three component RDS with a biologically implausible non-local term. Our computational method based on surface finite elements extends the method in [16] and is an alternative to the level set method considered in [7, 37]. The simulations illustrate that the model is capable of reproducing aspects of pseudopod-driven cell migration, described in [17], in both two and three space dimensions. We report on many widely used chemotaxis measures and observe values similar to experimental observations. We also note that the simulations exhibit a dilution effect at the tip of a pseudopod where the local maxima corresponding to an activator peak is reduced. This suggests experimental investigation of the relative importance of mechanical effects of membrane protrusion on the distribution of cell resident proteins is warranted.

We also investigated a model for the motion of fish keratocytes. The model appears to reproduce some experimental observations of the shapes of motile keratocyte cells and the experimental observation of the correlation between cell shape and speed [18]. The computational model in [18] reproduces the velocity-aspect ratio relationship. However, unlike our model, both polarisation and cell shapes are not explicitly modelled, with a parabolic actin profile at the leading edge assumed and the shape of the cell rear neglected. Studies [6] and [22] propose models where polarisation is modelled by equations in the bulk of the cell which are coupled to an evolution law for the cell surface. The import of our study is to show that a surface RDS coupled to a surface evolution law gives qualitatively similar results. A further contribution is the use of surface finite elements rather than the phase field method considered in [6] and [22]. This allows simulation of $3d$ keratocyte migration which studies [6] and [22] both note is computationally expensive with the phase field methodology. We do observe minor differences from [6], for example in the measures of deviation from reflection symmetry.

Our numerical experience suggests that some aspects of cell migration and chemotaxis can be captured by the Schnakenberg RDS (5.1). This RDS is considerably simpler from a mathematical analysis viewpoint than say the Meinhardt model (4.1). One can show that the model is well posed on evolving (planar) domains [48], which is an open question even on fixed domains for the Meinhardt model. As the two components are out of phase the model lends itself naturally to the case of a species that promotes protrusion (e.g., actin) and another that promotes retraction (e.g., myosin).

In this work it is our intention to present a framework for future modelling rather than suggest any definitive models for cell migration. We hope that future studies will employ the framework we have set out to refine existing models for cell motility and make predictions based on numerical simulations that can be used to direct and inform experimental studies.

ACKNOWLEDGMENTS

This research has been supported by the UK Engineering and Physical Sciences Research Council (EPSRC), Grant EP/G010404.

REFERENCES

- [1] D. Bray. *Cell movements: from molecules to motility*. Routledge, 2001.
- [2] A. Mogilner. Mathematics of cell motility: have we got its number? *Journal of mathematical biology*, 58(1):105–134, 2009.
- [3] A. Jilkin and L. Edelstein-Keshet. A comparison of mathematical models for polarization of single eukaryotic cells in response to guided cues. *PLoS Computational Biology*, 7(4), 2011.
- [4] J.C. Del Alamo, R. Meili, B. Alonso-Latorre, J. Rodríguez-Rodríguez, A. Aliseda, R.A. Firtel, and J.C. Lasheras. Spatio-temporal analysis of eukaryotic cell motility by improved force cytometry. *Proceedings of the National Academy of Sciences*, 104(33):13343, 2007.
- [5] M.L. Lombardi, D.A. Knecht, M. Dembo, and J. Lee. Traction force microscopy in dictyostelium reveals distinct roles for myosin ii motor and actin-crosslinking activity in polarized cell movement. *Journal of Cell Science*, 120(9):1624–1634, 2007.
- [6] F. Ziebert, S. Swaminathan, and I.S. Aranson. Model for self-polarization and motility of keratocyte fragments. *Journal of The Royal Society Interface*, 2011.
- [7] M.P. Neilson, J. Mackenzie, S. Webb, and R.H. Insall. Modelling cell movement and chemotaxis pseudopod based feedback. *SIAM Journal on Scientific Computing*, 33(3), 2011.
- [8] JM Oliver, JR King, KJ McKinlay, PD Brown, DM Grant, CA Scotchford, and JV Wood. Thin-film theories for two-phase reactive flow models of active cell motion. *Mathematical Medicine and Biology*, 22(1):53–98, 2005.
- [9] W. Alt and M. Dembo. Cytoplasm dynamics and cell motion: two-phase flow models. *Mathematical biosciences*, 156(1-2):207–228, 1999.
- [10] G. Dziuk and C.M. Elliott. Finite elements on evolving surfaces. *IMA journal of numerical analysis*, 27(2):262, 2007.
- [11] J.W. Barrett, H. Garcke, and R. Nürnberg. Parametric approximation of Willmore flow and related geometric evolution equations. *SIAM Journal on Scientific Computing*, 31:225, 2008.
- [12] L. Yang, J. Effler, B. Kutscher, S. Sullivan, D. Robinson, and P. Iglesias. Modeling cellular deformations using the level set formalism. *BMC systems biology*, 2(1):68, 2008.
- [13] C. Landsberg and A. Voigt. A multigrid finite element method for reaction-diffusion systems on surfaces. *Computing and Visualization in Science*, 13(4):177–185, 2010.
- [14] G. Dziuk and C.M. Elliott. Eulerian finite element method for parabolic pdes on implicit surfaces. *Interfaces and Free Boundaries*, 10(119-138):464, 2008.
- [15] R.I. Saye and J.A. Sethian. The voronoi implicit interface method for computing multiphase physics. *Proceedings of the National Academy of Sciences*, 108(49):19498–19503, 2011.
- [16] M.P. Neilson, J.A. Mackenzie, S.D. Webb, and R.H. Insall. Use of the parameterised finite element method to robustly and efficiently evolve the edge of a moving cell. *Integr. Biol.*, 2010.
- [17] R.H. Insall. Understanding eukaryotic chemotaxis: a pseudopod-centred view. *Nature Reviews Molecular Cell Biology*, 11(6):453–458, 2010.
- [18] K. Keren, Z. Pincus, G.M. Allen, E.L. Barnhart, G. Marriott, A. Mogilner, and J.A. Theriot. Mechanism of shape determination in motile cells. *Nature*, 453(7194):475–480, 2008.
- [19] J.D. Murray. *Mathematical biology*. Springer Verlag, 2003.
- [20] C. Venkataraman, T. Sekimura, E.A. Gaffney, P.K. Maini, and A. Madzvamuse. Modeling parr-mark pattern formation during the early development of amago trout. *Phys. Rev. E*, 84:041923, Oct 2011. doi: 10.1103/PhysRevE.84.041923. URL <http://link.aps.org/doi/10.1103/PhysRevE.84.041923>.

- [21] R. Barreira, C.M. Elliott, and A. Madzvamuse. The surface finite element method for pattern formation on evolving biological surfaces. *Journal of Mathematical Biology*, pages 1–25, 2011. ISSN 0303-6812.
- [22] D. Shao, W.J. Rappel, and H. Levine. Computational model for cell morphodynamics. *Physical review letters*, 105(10):108104, 2010.
- [23] David Traynor and Robert R. Kay. Possible roles of the endocytic cycle in cell motility. *Journal of Cell Science*, 120(14):2318–2327, 2007. doi: 10.1242/jcs.007732. URL <http://jcs.biologists.org/content/120/14/2318.abstract>.
- [24] K. Larripa and A. Mogilner. Transport of a 1d viscoelastic actin-myosin strip of gel as a model of a crawling cell. *Physica A: Statistical Mechanics and its Applications*, 372(1):113–123, 2006.
- [25] K. Deckelnick, G. Dziuk, and C.M. Elliott. Computation of geometric partial differential equations and mean curvature flow. *Acta Numerica*, 14:139–232, 2005. ISSN 0962-4929.
- [26] W. Helfrich. Elastic properties of lipid bilayers: theory and possible experiments. *Z. Naturforsch*, 28(11):693–703, 1973.
- [27] T.J. Willmore. *Riemannian geometry*. Oxford University Press, USA, 1997.
- [28] G. Dziuk. An algorithm for evolutionary surfaces. *Numerische Mathematik*, 58(1):603–611, 1990.
- [29] G. Dziuk and C.M. Elliott. L2-estimates for the evolving surface finite element method. *Math Comp (to appear)*, 2012.
- [30] C.M. Elliott and B. Stinner. Modeling and computation of two phase geometric biomembranes using surface finite elements. *Journal of Computational Physics*, 229(18):6585–6612, 2010.
- [31] C.M. Elliott and B. Stinner. Computation of two-phase biomembranes with phase dependent material parameters using surface finite elements. *CiCP (to appear)*, 2012.
- [32] CJ Heine. Isoparametric finite element approximation of curvature on hypersurfaces. Preprint.
- [33] G. Dziuk. Computational parametric Willmore flow. *Numerische Mathematik*, 111(1):55–80, 2008.
- [34] A. Schmidt and K.G. Siebert. *Design of adaptive finite element software: The finite element toolbox ALBERTA*. Springer Verlag, 2005.
- [35] Timothy A. Davis. Algorithm 832: Umfpack v4.3—an unsymmetric-pattern multifrontal method. *ACM Trans. Math. Softw.*, 30:196–199, June 2004. ISSN 0098-3500. doi: <http://doi.acm.org/10.1145/992200.992206>. URL <http://doi.acm.org/10.1145/992200.992206>.
- [36] H. Meinhardt. Orientation of chemotactic cells and growth cones: models and mechanisms. *Journal of Cell Science*, 112(17):2867, 1999.
- [37] M.P. Neilson, D.M. Veltman, P.J.M. van Haastert, S.D. Webb, J.A. Mackenzie, and R.H. Insall. Chemotaxis: A feedback-based computational model robustly predicts multiple aspects of real cell behaviour. *PLoS biology*, 9(5):e1000618, 2011.
- [38] D.P. Amarasinghe, A. Aylwin, P. Madhavan, and C. Pettitt. Biomembranes report. MASDOC, RSG, 2011.
- [39] I. Hecht, M.L. Skoge, P.G. Charest, E. Ben-Jacob, R.A. Firtel, W.F. Loomis, H. Levine, and W.J. Rappel. Activated membrane patches guide chemotactic cell motility. *PLoS Computational Biology*, 7(6):e1002044, 2011.
- [40] S.Y. Cheng, S. Heilman, M. Wasserman, S. Archer, M.L. Shuler, and M. Wu. A hydrogel-based microfluidic device for the studies of directed cell migration. *Lab Chip*, 7(6):763–769, 2007.
- [41] WS Ramsey. Analysis of individual leucocyte behavior during chemotaxis. *Experimental cell research*, 70(1):129–139, 1972.
- [42] L. Bosgraaf and P.J.M. Van Haastert. Navigation of chemotactic cells by parallel signaling to pseudopod persistence and orientation. *PloS one*, 4(8):e6842, 2009.
- [43] I. Hecht, H. Levine, W.J. Rappel, and E. Ben-Jacob. “self-assisted” amoeboid navigation in complex environments. *PloS one*, 6(8):e21955, 2011.
- [44] R. Grima. Directed cell migration in the presence of obstacles. *Theoretical Biology and Medical Modelling*, 4(1):2, 2007.

- [45] L. Bosgraaf, P.J.M. van Haastert, and T. Bretschneider. Analysis of cell movement by simultaneous quantification of local membrane displacement and fluorescent intensities using quimp2. *Cell motility and the cytoskeleton*, 66(3):156–165, 2009.
- [46] T.D. Pollard and G.G. Borisy. Cellular motility driven by assembly and disassembly of actin filaments. *Cell*, 112(4):453–465, 2003.
- [47] R. Lefever and I. Prigogine. Symmetry-breaking instabilities in dissipative systems II. *J. chem. Phys.*, 48:1695–1700, 1968.
- [48] Chandrasekhar Venkataraman, Omar Lakkis, and Anotida Madzvamuse. Global existence for semi-linear reaction–diffusion systems on evolving domains. *Journal of Mathematical Biology*, 64:41–67, 2012. ISSN 0303-6812. URL <http://dx.doi.org/10.1007/s00285-011-0404-x>. 10.1007/s00285-011-0404-x.